

## References

- ANDERSON, A. M., D. A. HAUKOS & J. T. ANDERSON (1999): Diet composition of three anurans from the playa wetlands of northwest Texas. — *Copeia* 1999: 515–520.
- HOUSTON, W. W. K. (1973): The food of the common frog *Rana temporaria* on high moorland in northern England. — *Journal of Zoology*, London 171: 153–165.
- ITÄMIES, J. & P. KOSKELA (1970). On the diet of the common frog (*Rana temporaria L.*). — *Aquilo Ser. Zool.* 10: 53–60.
- JENSSEN, T. A. (1967): Food habits of the green frog, *Rana clamitans*, before and during metamorphosis. — *Copeia* 1967: 214–218.
- KUZMIN, S. L. (1984): Age changes of feeding in *Hynobius keyserlingii*. — *Zoological Zhurnal*, Moskau 63: 1055–1060 (in Russian with English abstract).
- KUZMIN, S. L. (1997): Feeding of amphibians during metamorphosis. — *Amphibia-Reptilia*, Leiden 18: 121–131.
- LOMAN, J. (1979): Food, feeding rates and prey-size selection in juvenile and adult frogs *Rana arvalis NILS.* and *R. temporaria L.* — *Ekol. pol.* 27: 581–601.

## Fehlpaarung zwischen *Rana temporaria* und *Salamandra salamandra* mit tödlichem Ausgang

MICHAEL LINNENBACH

(Schubertstr. 9, D-69226 Nussloch)

### Misguided copulation of *Rana temporaria* and *Salamandra salamandra* proved fatal.

At the 28.3.2000 I observed a male *Rana temporaria* clasping a female *Salamandra salamandra* to death.

**Key words:** *Rana temporaria*, *Salamandra salamandra*, misguided copulation.